arXiv:1301.7639v2 [quant-ph] 26 Feb 2013

On the real matrix representation of PT-symmetric operators

Francisco M. Fernández

INIFTA (UNLP, CCT La Plata-CONICET), Blvd. 113 y 64 S/N, Sucursal 4, Casilla de Correo 16, 1900 La Plata, Argentina

E-mail: `fernande@quimica.unlp.edu.ar`

Abstract. We discuss the construction of real matrix representations of PT-symmetric operators. We show the limitation of a general recipe presented some time ago for non-Hermitian Hamiltonians with antiunitary symmetry and propose a way to overcome it. Our results agree with earlier ones for a particular case.

It is well known that some complex-valued non-Hermitian operators exhibit real eigenvalues [1] (and references therein). In order to provide a simple and general explanation of this fact Bender et al [2] showed that it is possible to obtain a basis set so that the matrix representation of such operators is real. The argument is based on the existence of an antiunitary symmetry.

An antiunitary operator satisfies

$$\left\langle \hat{A}\phi \middle| \hat{A}\psi \right\rangle = \left\langle \phi \right| \psi\rangle^{*} \tag{1}$$

for arbitrary states $|\phi\rangle$ and $|\psi\rangle$. It can be written as the product of a unitary operator $\hat{U}$ and the complex conjugation operator $\hat{K}$: $\hat{A} = \hat{U}\hat{K}$. Bender et al. [2] considered Hamiltonians $\hat{H}$ with antiunitary symmetry $[\hat{H}, \hat{A}] = 0$, where $\hat{A}$ satisfies the additional condition

$$\hat{A}^{2k} = \hat{1},\ k \text{ odd} \tag{2}$$

They proved that for an $\hat{A}$-adapted basis set $\{|n_A\rangle\}$

$$\hat{A}^{k} |n_A\rangle = |n_A\rangle \tag{3}$$

the matrix elements of the operator are real

$$\langle m_A| \hat{H} |n_A\rangle = \langle m_A| \hat{H} |n_A\rangle^{*} \tag{4}$$

Those authors proposed to construct $|n_A\rangle$ as

$$|n_A\rangle = |n\rangle + \hat{A}^{k} |n\rangle \tag{5}$$

where $\{|n\rangle\}$ is any orthonormal basis set.

It is not difficult to prove that this recipe does not apply to any basis set. According to equation (2) we can find a basis set $\{|n, \sigma\rangle\}$ that satisfies

$$\hat{A}^{k} |n, \sigma\rangle = \sigma |n, \sigma\rangle\,,\ \sigma = \pm 1 \tag{6}$$

Consequently, all the vectors

$$\left|n,\sigma\right\rangle_A = \left|n,\sigma\right\rangle + \hat{A}^k\left|n,\sigma\right\rangle = (1+\sigma)\left|n,\sigma\right\rangle \tag{7}$$

with $\sigma = -1$ vanish and the resulting $\hat{A}$-adapted vector set is not complete. We conclude that the basis set $\{|n\rangle\}$ should be chosen carefully in order to apply the recipe of Bender et al [2].

We can construct the basis set $\{|n,\sigma\rangle\}$ from any orthonormal basis set $\{|n\rangle\}$ in the following way

$$\left|n,\sigma\right\rangle = N_{n,\sigma}\hat{Q}_\sigma\left|n\right\rangle,\ \hat{Q}_\sigma = \frac{1}{2}\left(1+\sigma\hat{A}^k\right) \tag{8}$$

where $N_{n,\sigma}$ is a suitable normalization factor. It already satisfies equation (6) because $\hat{A}^k\hat{Q}_\sigma = \sigma\hat{Q}_\sigma$. In order to overcome the shortcoming in the recipe of Bender et al [2] we define the $\hat{A}$-adapted basis set $B_A = \{\left|n_A^+\right\rangle = \left|n,1\right\rangle, \left|n_A^-\right\rangle = i\left|n,-1\right\rangle\}$. Note that the vectors $\left|n_A^\pm\right\rangle$ satisfy the requirement (3) and that B_A is complete. In principle there is no guarantee of orthogonality, but such a difficulty does not arise in the examples discussed below.

The vectors $|v_n^\pm\rangle = \frac{1}{\sqrt{2}}\left(\left|n_A^+\right\rangle \pm \left|n_A^-\right\rangle\right)$ also satisfy the requirement (3) and in the particular case of a two-dimensional space they lead to the $\hat{A}$-adapted basis set chosen by Bender et al [2] to introduce the issue by means of a simple example .

As another example consider the parity-time antiunitary operator $\hat{A} = \hat{P}\hat{T}$ that is a particular case of the equations above with $k = 1$. Let $\{|n\rangle, n = 0,1,\ldots\}$ be the basis set of eigenvectors of the Harmonic oscillator $\hat{H}_0 = \hat{p}^2 + \hat{x}^2$ that are real and satisfy $\hat{P}\left|n\right\rangle = (-1)^n\left|n\right\rangle$ so that $\hat{A}\left|n\right\rangle = (-1)^n\left|n\right\rangle$. It is clear that the recipe proposed by Bender et al [2] does not apply to this simple case. On the other hand, present recipe yields the $\hat{A}$-adapted basis set $B_A^{HO} = \{\left|2n\right\rangle, i\left|2n+1\right\rangle, n = 0,1,\ldots\}$ which is obviously complete.

Every vector of the orthonormal basis set B_A^{HO} in the coordinate representation can be expressed as a linear combination of the elements of the nonorthogonal basis set $\{f_n(x) = e^{-x^2/2}(ix)^n, n = 0,1,\ldots\}$. By means of a slight generalization of the latter

Znojil [3] derived a recurrence relation with real coefficients for a family of complex anharmonic potentials. This author also constructed a real matrix representation of a PT-symmetric oscillator in terms of the eigenvectors of $\hat{A} = \hat{P}\hat{T}$ [4]. Note that his vectors $|S_n\rangle$ and $|L_n\rangle$ are our $|n, 1\rangle$ and $|n, -1\rangle$ respectively.

[1] Bender C M 2007 *Rep. Prog. Phys.* **70** 947.

[2] Bender C M, Berry M V, and Mandilara A 2002 *J. Phys. A* **35** L467.

[3] Znojil M 1999 *J. Phys. A* **32** 7419.

[4] Znojil M 2002 *J. Nonlin. Math. Phys.* **9** 122.